



VU Research Portal

A re-discovered treasure: database of mycorrhizal intensity for 3000 vascular plant species across multiple sites throughout former Soviet Union territory.

Akmetzhanova, A.A.; Soudzilovskaia, N.A.; Onipchenko, V.G.; Cornwell, W.K.; Agafonov, V.; Selivanov, I.A.; Cornelissen, J.H.C.

published in

Ecology

2012

DOI (link to publisher)

[10.1890/11-1749.1](https://doi.org/10.1890/11-1749.1)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Akmetzhanova, A. A., Soudzilovskaia, N. A., Onipchenko, V. G., Cornwell, W. K., Agafonov, V., Selivanov, I. A., & Cornelissen, J. H. C. (2012). A re-discovered treasure: database of mycorrhizal intensity for 3000 vascular plant species across multiple sites throughout former Soviet Union territory. *Ecology*, 93, 689-690. <https://doi.org/10.1890/11-1749.1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

A rediscovered treasure: mycorrhizal intensity database for 3000 vascular plant species across the former Soviet Union

Ecological Archives E093-059

ASEM A. AKHMETZHANOVA,^{1,3} NADEJDA A. SOUDZILOVSKAIA,^{2,5} VLADIMIR G. ONIPCHENKO,^{1,3} WILL K. CORNWELL,¹
VLADIMIR A. AGAFONOV,³ IVAN A. SELIVANOV,^{4,6} AND JOHANNES H. C. CORNELISSEN²

¹*Department of Geobotany, Faculty of Biology, Moscow State University, RU-119991 Moscow, Russia*

²*Systems Ecology, Department of Ecological Science, Faculty of Earth and Life Sciences, VU University, De Boelelaan 1085, NL-1081HV Amsterdam, The Netherlands*

³*Teberda State Biosphere Reserve, Teberda, Badukskii 1, 369210, Karachai-Cherkessian Republik, Russia*

⁴*Perm State Pedagogic University, Ulitsa Sibirskaya, 24, Perm 614990 Russia*

Abstract. The symbiosis between vascular plants and mycorrhizal fungi is paramount for carbon and nutrient cycling in most of the world's ecosystems. Most vascular plant species are associated with mycorrhizal fungal partners, and the association is essential for the carbon and nutrition economies of both partners. However, despite its clear importance, data on this symbiosis are lacking: for most vascular plant species, mycorrhizal type is unknown. Very rarely is there data on the levels of mycorrhizal infection intensity in multiple habitats.

We translated and digitized a huge data set on vascular plant mycorrhizal intensity throughout the former Soviet Union, previously available only as a hard copy appendix of the doctoral thesis of Ivan A. Selivanov published in Russian in 1976 and not accessible to the international research community. We updated the taxonomic plant nomenclature to the International Plant Name Index and adjusted mycorrhizal and ecological terminology according to the modern international literature.

The database contains 7445 records on mycorrhizal infection type and intensity of 2970 plant species from 155 families, in 154 sites, situated across the former Soviet Union (mostly on the territory of the current Russia, Ukraine, and Kazakhstan), comprising together extensive geological, topographic, and climatic gradients. The data set includes percentage infection for each species–site combination for arbuscular, ericoid, arbutoid, endo-mycorrhizal, dark septate, orchid- and ecto-mycorrhizal fungi. Each record has a detailed description of geography. For many records, soils and host plant community are described. Most of the sites are natural; 10 sites are situated in botanical gardens. For 1291 species the intensity of mycorrhizal infection is quantified in multiple plant communities (2–57). The remaining species are described at single sites.

Selivanov developed his own methods for quantifying mycorrhizal infection intensity. These methods are comparable, but not identical to, the methodology commonly used today. Based on our own sampling of 99 plant species collected in two distant sites (Caucasus [Russia] and Abisko [Sweden]), we provide a simple equation for data conversion between the two methods.

The availability of this database will help to provide answers to important questions concerning biogeochemical cycling, climate change impacts, and co-evolution of plants and fungi.

Key words: *arbuscular mycorrhiza; carbon cycling; ecto-mycorrhiza; ericoid mycorrhiza; Kazakhstan; mycorrhizal infection; mycorrhizal type; nutrient cycling; root colonization; Russia; symbiosis; Ukraine.*

The complete data sets corresponding to abstracts published in the Data Papers section of the journal are published electronically in *Ecological Archives* at (<http://esapubs.org/archive>). (The accession number for each Data Paper is given directly beneath the title.)

Manuscript received 26 September 2011; revised 29 November 2011; accepted 30 November 2011. Corresponding Editor: W. K. Michener.

⁵ Corresponding author. E-mail: nadia.soudzilovskaia@falw.vu.nl

⁶ Deceased.